

INFORMATION ABOUT ADVERSE EVENTS FOLLOWING VACCINATION AND WHAT TO EXPECT (영어)

COMMON MILD SIDE EFFECTS AND WHAT YOU CAN DO TO HELP:

- If you have swelling or pain after vaccination, you can put a [clean dry towel on the affected area](#) and [put a cool, damp washcloth on top of it](#).
- If you have a mild fever after vaccination, [drink plenty of water and get sufficient rest](#).
- If you experience discomfort due to fever or muscle pain after vaccination, [taking fever medicine or pain relievers](#) can help.
- * You can prepare an acetaminophen-based fever or pain medication in advance, but take them only if you have fever or body ache after vaccination.

SEE A DOCTOR IF YOU EXPERIENCE THE FOLLOWING:

- [Swelling, pain, or redness](#) of the vaccination spot that does not improve after 48 hours
- Symptoms such as [shortness of breath, chest pain, persistent abdominal pain, and leg swelling](#) within 4 weeks after vaccination
- Severe or persistent headache that lasts more than 2 days after vaccination and does [not respond to pain medications, is not controlled, or accompanied with blurred vision](#)
- [Sudden loss of energy or other unusual symptoms](#) after vaccination
- [Bruises or bleeding](#) not occurring in parts of your body other than the spot where you got the vaccine shot

DIAL 119 OR GO TO AN EMERGENCY ROOM IF YOU EXPERIENCE THE FOLLOWING:

- Severe [breathing difficulties](#) or severe [dizziness](#) after vaccination
- [Swelling of lips or face](#) or [severe hives](#) all over the body after the vaccination
- Sudden [loss of consciousness or fainting](#)

<Translated by Danuri Call Center 1577-1366>